

Colour Change Measurements of Gravitational Microlensing Events by Using the Difference Image Analysis Method

Cheongho Han and Seong-Hong Park

*Department of Astronomy & Space Science,
Chungbuk National University,
Chongju, Korea 361-763,
cheongho,parksh@astro.chungbuk.ac.kr*

Accepted:

Received:

ABSTRACT

Detecting colour changes of a gravitational microlensing event induced by the limb-darkened extended source effect is important to obtain useful information both about the lens and source star. However, precise measurements of the colour changes are hampered by blending, which also causes colour changes of the event. In this paper, we show that although the colour change measured from the subtracted image by using the recently developed photometric method of the “difference image analysis” (DIA) differs from the colour change measured by using the conventional method based on the extraction of the individual source stars’ point spread functions, the curve of the colour changes (colour curve) constructed by using the DIA method enables one to obtain the same information about the lens and source star, but with significantly reduced uncertainties due to the absence of blending. We investigate the patterns of the DIA colour curves for both single lens and binary lens events by constructing colour change maps.

Key words: gravitational lensing – stars: fundamental parameters (colours) – technique: photometric

1 INTRODUCTION

The amplification of source flux caused by gravitational lensing can become theoretically infinite. Points in the source plane on which the amplification of a point source becomes infinite are called caustics. For a point lens, the caustic is a single point behind the lens. For a binary lens, the number of caustics is multiple and they form closed curves. For a real microlensing event, however, the source star has an extended size, and thus the observed amplification deviates from that of a point source event and becomes always finite: extended source effect (Schneider, Ehlers, & Falco 1992; Witt & Mao 1994). The deviation of the light curve caused by the extended source effect becomes important as the source passes very close to the lens caustics.

Detection of the extended source effect is important because one can obtain useful information both about the lens and source star. First, a caustic crossing event provides an opportunity to measure how long it takes for the caustics to transit the face of the source star. By using the source radius crossing time t_* , along with an independent determination of the source star size θ_* , one can determine the lens proper motion with respect to the source star by $\mu = \theta_*/t_*$ (Witt & Mao 1994; Gould 1994; Maoz & Gould 1994; Nemiroff &

Wickramasinghe 1994; Loeb & Sasselov 1995; Alcock et al. 1997b, 1997c, 1999a; Afonso et al. 1998; Udalski et al. 1998; Albrow et al. 1999a). Once the value of μ is determined, the mass and location of the lens can be significantly better determined. Second, by analyzing the light curve of an event with the source trajectory passing very close to the caustics, one can investigate the surface structure of the source star such as the surface intensity profile and spots (Valls-Gabaud 1994, 1998; Sasselov 1997; Gaudi & Gould 1999; Heyrovský & Sasselov 2000; Albrow et al. 1999b; Han et al. 2000).

In addition to detecting deviations in the light curve, the extended source effect can also be detected by measuring the colour changes of the source star during the event. The colour changes occur due to the differential amplification over the limb-darkened source star surface (see § 3.1). Throughout this paper, we use a term ‘colour curve’ to refer to the colour changes of an event induced by the extended source effect as a function of time. Detection of the extended source effect from the colour measurements has the following advantages compared to the detection from a single band photometry. First, by detecting colour changes one can increase the chance to detect the extended source effect. If the source star approaches very close to the lens caustics but does not transit the caustics, the amplification induced by

the extended source effect simply masquerades as changes in lensing parameters, and thus cannot be detected. By contrast, the colour change cannot be mimicked by the change in lensing parameters, because a point source lensing event should be achromatic (Gould & Welch 1996). Second, by measuring the colour changes one can determine the lens proper motion with relative ease. This is because one can measure t_* , and thus μ , by simply measuring the turning time of the colour curve (Han, Park, & Jeong 2000, see § 4 for more details). To determine t_* from the light curve, on the other hand, it is required to fit the overall light curve.

However, precise measurements of the colour changes induced by the extended source effect is hampered by blending, which also causes colour changes of the event (Kamionkowski 1995; Buchalter, Kamionkowski, & Rich 1996). To increase the event rate, the current lensing experiments are being conducted towards very dense star fields such as the Galactic bulge and the Magellanic Clouds (Alcock et al. 1993; Aubourg et al. 1993; Udalski et al. 1993; Alard & Guibert 1997). When the brightness of a source star located towards these very dense star fields is measured by using the conventional method based on the extraction of the individual source stars' point spread functions, it is very likely that the measured source star flux is affected by the unwanted light from nearby unresolved stars (Di Stefano & Esin 1995; Alard 1997; Alcock et al. 1997a; Palanque-DeLabrouille et al. 1998; Woźniak & Paczyński 1997; Han 1999; Han, Jeong, & Kim 1998). Since blended stars in general have different colours from the colour of the lensed source star, the measured colour is affected by blending. Han, Park, & Jeong (2000) pointed out the seriousness of blending in colour change measurements by demonstrating that even for a very small fraction ($\lesssim 2\%$) of blended light, the colour change caused by blending can be equivalent to the colour change induced by the extended source effect. Therefore, for the precise measurements of the colour changes induced by the extended source effect, it will be essential to correct for or remove the blending effect.

Although numerous methods have been proposed for the correction of blending (Alard, Mao, & Guibert 1995; Alard 1996; Han 1997, 1998; Goldberg 1998; Goldberg & Woźniak 1998; Han & Kim 1999), most of these methods either have limited applicabilities only to several special cases of blended events or are less practical due to the requirement of space observations.* On the other hand, with the recently developed photometric technique of the “difference image analysis” (DIA) method, one can measure the blending-free light variations of general microlensing events (Tomaney & Crofts 1996; Alard & Lupton 1998; Alard 1999; Alcock et al. 1999b, 1999c). The DIA method detects and measures the variation of the source star flux by subtracting an observed image from a convolved and normalized reference image. Since the blended light is subtracted during the image subtraction process, one can improve the photometric precision by removing the effect of blending. At the moment

* We note, however, that the MACHO group had *Hubble Space Telescope* followup observations of the source star for one of the events they have detected to clearly identify the lensed source star. From these observations, they could accurately estimate the lens mass by correcting blending (NASA press release #STScI-PR00-03).

most microlensing experiments do not use the DIA method in their data analysis. Using the DIA method would require that either the whole microlensing survey process should switch to the DIA method, or that systematic re-processing of the light curves should be performed once events have been detected using the conventional photometric method.

Due to the methodological difference of the DIA photometry from the conventional photometry, the colour change of an event measured by the using the DIA method (DIA colour change) differs from that measured by the conventional method (see § 3). In this paper, we show, however, that the DIA colour curve of an event enables one to obtain the same information about the lens and source star, but with significantly reduced uncertainties due to the absence of blending. We investigate the patterns of the DIA colour curves for both single and binary lens events by constructing colour change maps.

2 GRAVITATIONAL AMPLIFICATION

2.1 Point Source Events

If a microlensing event with a point source is caused by a single lens, the light curve of the event is represented by

$$A_p = \frac{u^2 + 2}{u(u^2 + 4)^{1/2}}; \quad u = \left[\left(\frac{t - t_0}{t_E} \right)^2 + \beta^2 \right]^{1/2}, \quad (1)$$

where the subscript ‘p’ denotes the event is occurred to a point source, u represents the lens-source separation normalized by the angular Einstein ring radius θ_E , and the lensing parameters β , t_0 , and t_E represent the impact parameter of the lens-source encounter, the time of maximum amplification, and the Einstein ring radius crossing time (Einstein time scale), respectively. The angular Einstein ring represents the effective region of gravitational amplification and its size is related to the physical parameters of the lens by

$$\theta_E = \left(\frac{4GM}{c^2} \frac{D_{ls}}{D_{ol}D_{os}} \right)^{1/2}, \quad (2)$$

where M is the mass of the lens and D_{ol} , D_{ls} , and D_{os} represent the separations between the observer, the lens, and the source, respectively. When the source is located within the Einstein ring, the source star amplification becomes greater than $3/\sqrt{5}$.

If an event is caused by a binary lens, on the other hand, the resulting light curve differs from that of a single lens event. When lengths are normalized by the combined angular Einstein ring radius, which represents the angular Einstein ring radius with a lens mass equal to the total mass of the binary system, the lens equation in complex notation for the binary lens system is represented by

$$\zeta = z + \frac{m_1}{\bar{z}_1 - \bar{z}} + \frac{m_2}{\bar{z}_2 - \bar{z}}, \quad (3)$$

where m_1 and m_2 are the mass fractions of individual lenses (and thus $m_1 + m_2 = 1$), z_1 and z_2 are the positions of the lenses, $\zeta = \xi + i\eta$ and $z = x + iy$ are the positions of the source and images, and $\bar{z}$ denotes the complex conjugate of z (Witt 1990). The amplification of each image, $A_{p,i}$, is given by the Jacobian of the transformation (3) evaluated at the images position, i.e.

$$A_{p,i} = \left(\frac{1}{|\det J|} \right)_{z=z_i}; \quad \det J = 1 - \frac{\partial \zeta}{\partial \bar{z}} \frac{\partial \bar{\zeta}}{\partial z}. \quad (4)$$

Then the total amplification of the event is given by the sum of the individual amplifications, i.e. $A_p = \sum_i A_{p,i}$. The caustic refers to the source position on which the total amplification becomes infinite, i.e. $\det J = 0$.

2.2 Extended Source Events

If the source of an event has an extended size, the light curve of the event deviates from that of a point source event. For both single and binary lens events, the amplification of an extended source event is the weighted mean of the amplification factor over the source star disk, i.e.

$$A_\nu = \frac{\int_0^{r_*} I_\nu(r) A_p(|\mathbf{r} - \mathbf{r}_L|) r dr}{\int_0^{r_*} I_\nu(r) r dr}, \quad (5)$$

where $I_\nu(r)$ is the radial surface intensity distribution of the source star with a radius r_* and the vector $\mathbf{r}_L$ and $\mathbf{r}$ represent the displacement vector of the source star center with respect to the lens and the orientation vector of a point on the source star surface with respect to the center of the source star, respectively.

3 COLOUR CHANGES

3.1 Measured by the Conventional Photometric Method

In addition to causing deviations in the light curve from that of a point source event, the extended source effect makes the gravitational amplification become wavelength dependent, causing colour changes during the event. The colour changes occur due to the wavelength dependency of the source star radial surface intensity profile caused by limb darkening. If we define $F_{0,\nu i} = 2\pi \int_0^{r_*} I_{\nu i}(r) r dr$, where $i = 1, 2$, and $m_{\nu i}$ as the unamplified source star fluxes and the corresponding magnitudes measured at two different wavelength passbands $\nu 1$ and $\nu 2$, the colour of the *un-blended* source star measured at a time t by using the conventional method is computed by

$$(m_{\nu 2} - m_{\nu 1})_0(t) = -2.5 \log \left[\frac{A_{\nu 2}(t) F_{0,\nu 2}}{A_{\nu 1}(t) F_{0,\nu 1}} \right]. \quad (6)$$

Then the colour curve of the event is represented by

$$\Delta(m_{\nu 2} - m_{\nu 1})_0(t) = -2.5 \log \left\{ \left[\frac{A_{\nu 2}(t)}{A_{\nu 1}(t)} \right] \left[\frac{A_{\nu 2}(t_{\text{ref}})}{A_{\nu 1}(t_{\text{ref}})} \right]^{-1} \right\}, \quad (7)$$

where t_{ref} represents the reference time for the colour change measurements. For a point source event, $A_{\nu 1}(t) = A_{\nu 2}(t)$ (i.e. achromatic), and thus $(m_{\nu 2} - m_{\nu 1})_0 = -2.5 \log(F_{0,\nu 2}/F_{0,\nu 1})$, implying that the colour of the source during the event equals to that of the un-amplified source. For a limb-darkened extended source event, on the other hand, as the source passes close to caustics, different parts of the source star disk with varying surface intensity and spectral energy distribution are amplified by different amount

due to the differences in distance to the lens. As a result, the amplification becomes wavelength dependent, i.e. $A_{\nu 1}(t) \neq A_{\nu 2}(t)$, and the colour changes during the event. Once the colour curve of the event is constructed, the lens proper motion and the source star surface intensity profile are determined by statistically comparing the observed colour curve to the theoretical curves with various models of limb darkening and source star size.

However, precise measurement of the colour changes induced by the extended source effect is hampered by blending, which is another mechanism causing colour changes of microlensing events. If one defines $B_{\nu i}$ as the blended amount of flux in two passbands, the measured colour of a blended event becomes

$$\begin{aligned} (m_{\nu 2} - m_{\nu 1})(t) &= -2.5 \log \left[\frac{A_{\nu 2}(t) F_{0,\nu 2} + B_{\nu 2}}{A_{\nu 1}(t) F_{0,\nu 1} + B_{\nu 1}} \right] \\ &= -2.5 \log \left[\frac{A_{\nu 2}(t) + f_{\nu 2}}{A_{\nu 1}(t) + f_{\nu 1}} \right], \end{aligned} \quad (8)$$

where $f_{\nu i} = B_{\nu i}/F_{0,\nu i}$ represent the ratios between the blended and the baseline source flux in the individual passbands. Then the colour curve of a *blended* event is represented by

$$\begin{aligned} \Delta(m_{\nu 2} - m_{\nu 1}) &= \\ &-2.5 \log \left\{ \left[\frac{A_{\nu 2}(t) + f_{\nu 2}}{A_{\nu 1}(t) + f_{\nu 1}} \right] \left[\frac{A_{\nu 2}(t_{\text{ref}}) + f_{\nu 2}}{A_{\nu 1}(t_{\text{ref}}) + f_{\nu 1}} \right]^{-1} \right\}. \end{aligned} \quad (9)$$

One finds that equation (9) includes two additional blending parameters of $f_{\nu 1}$ and $f_{\nu 2}$ compared to the un-blended event colour curve in equation (7). Therefore, to determine the lens proper motion and the source star surface intensity profile from the fit of the blended event colour curve, it is required to include these additional parameters, causing increased uncertainties in the determined quantities.

3.2 Measured by the DIA Method

The colour changes induced by the extended source effect can also be measured by using the DIA method. The flux of a source star measured from the subtracted image by using the DIA method is represented by

$$F_\nu = F_{\nu,\text{obs}} - F_{\nu,\text{ref}} = (A_\nu - 1)F_{\nu,0}, \quad (10)$$

where $F_{\nu,\text{obs}} = A_\nu F_{0,\nu} + B_\nu$ and $F_{0,\text{ref}} = F_{0,\nu} + B_\nu$ represent the source star fluxes measured from the images obtained during the progress of the event and from the reference image, respectively. Then the DIA colour curve of the event is represented by

$$\begin{aligned} \Delta(m_{\nu 2} - m_{\nu 1})_{\text{DIA}} &= \\ &-2.5 \log \left\{ \left[\frac{A_{\nu 2}(t) - 1}{A_{\nu 1}(t) - 1} \right] \left[\frac{A_{\nu 2}(t_{\text{ref}}) - 1}{A_{\nu 1}(t_{\text{ref}}) - 1} \right]^{-1} \right\}. \end{aligned} \quad (11)$$

From equation (11), one finds that the DIA colour curve does not depend on the blending parameters, and thus it is free from blending. One also finds that although the DIA colour curve has a different form from the curve of the un-blended event constructed by using the conventional method [cf. equation (7)], both curves depend on the same parameters of $A_{\nu 1}$ and $A_{\nu 2}$. Therefore, from the DIA colour




Figure 1. The colour map (upper panel) and curves (lower panel) for a single lens system. The map is presented as a function of lens position (x_L, y_L) with respect to the source star (the thick solid circle). The source star has an angular radius (normalized by θ_E) of $\theta_* = 0.1$. The reference of the colour change is chosen so that $\Delta(B-I)_{\text{DIA}} = 0$ when the source star is not gravitationally amplified. Grey scale becomes brighter (darker) as the colour of the source star becomes redder (bluer), and the tone of the grey scale changes with a colour change interval of 0.02 mag from the darkest region where $\Delta(B-I)_{\text{DIA}} \leq -0.06$ mag to the brightest region where $\Delta(B-I)_{\text{DIA}} > 0.04$. The lens trajectories (the straight lines in the upper panel) and their corresponding colour curves are drawn by the same line types.

curve one can obtain the same information about the lens and source star as that obtained from the colour curve constructed by using the conventional method, but with significantly reduced uncertainties due to the absence of blending.

4 PATTERNS OF DIA COLOUR CHANGE CURVES

In this section, we investigate the patterns of the DIA colour curves for both single and binary lens events. To see the

patterns, we construct the colour maps, which represent the colour changes as a function of source (lens) position with respect to the position of the lens (source). The constructed maps are presented in the upper panel of Figure 1 for the single lens system and Figure 2 for the binary lens system. For the construction of the maps, we assume that the source star has an angular radius (normalized by θ_E) of $\theta_* = 0.1$ and it is observed in B and I bands. The un-blended colour of the source star before amplification is $(B-I)_0 = 2.15$ mag, which corresponds to that of a K-type giant (Allen 1973; Schmidt-Kaler 1982; Peletier 1989). For the surface intensity profile, we adopt a linear form of

$$I_\nu(r) = 1 - C_\nu \left[1 - \sqrt{1 - (r/r_*)^2} \right], \quad (12)$$

where the limb-darkening coefficients are $C_B = 0.912$ and $C_I = 0.053$, respectively, which correspond to those of a K giant with $T_{\text{eff}} = 4,750$ K, $\log g = 2.0$, and a metallicity similar to the sun (Van Hamme 1993). For the binary lens system, we adopt a binary separation (also normalized by θ_E) of $\ell = 1.0$ and a mass ratio of $q = m_1/m_2 = 1.0$. The single lens map is presented as a function of lens positions (x_L, y_L) with respect to the source star (the thick solid circle with its center at the origin), while the binary lens map is presented as a function of source position (ξ, η) with respect to the center of the binary system. All lengths in the maps are normalized by θ_E . For both maps, we choose the reference of the colour change measurements to be $\Delta(B-I)_{\text{DIA}} = 0$ when the source star is not gravitationally amplified. Grey scale is drawn so that it becomes brighter (darker) as the colour of the source star becomes redder (bluer), and the tone of the grey scale changes for every colour change of 0.02 mag. For the single lens map, the grey tone change from the darkest region where $\Delta(B-I)_{\text{DIA}} \leq -0.06$ mag to the brightest region where $\Delta(B-I)_{\text{DIA}} > 0.04$ mag. For the binary lens map, the tone changes from the darkest region where $\Delta(B-I)_{\text{DIA}} \leq -0.05$ mag to the brightest region where $\Delta(B-I)_{\text{DIA}} > 0.07$ mag. To better show the fine structures of the binary lens event map in the region near the caustics (marked by thick solid curves), the region enclosed by a box (drawn by a short-dashed line) is expanded and presented in the upper panel of Figure 3. The straight lines in the upper panels of Figure 1 and 3 represent various lens (for the single lens case) or source (for the binary lens case) trajectories and the colour curves resulting from the trajectories are presented in the lower panels of the individual figures. The line types of the colour curves are selected so that they match with those of the corresponding lens or source trajectories.

From the colour map of the single lens system and the resulting colour curves, one finds the following patterns. First, the iso-colour-change contours are concentric circles with their center at the center of the source star. Due to the radial symmetry of the map, all the resulting colour curves are symmetric with respect to the time of maximum amplification. Second, the colour does not monotonically change as a function of the separation between the lens and the center of the source star, $r_L = \sqrt{x_L^2 + y_L^2}$. Outside the source star disk, the colour becomes redder as r_L decreases, but within the disk it becomes bluer with the decreasing value of r_L . As a result, while the colour of the non-source transit event continues to become redder as the lens approaches the




Figure 2. The colour map for a binary lens system. The map is presented as a function of source star position (ξ, η) with respect to the positions of the binary lenses (marked by ‘x’). The source star has an angular radius $\theta_* = 0.1$. The binary system has a separation (also normalized by θ_E) between its components of $\ell = 1.0$ and a mass ratio of $q = 1.0$. The closed figure drawn by a thick solid curve represents the caustics. Grey scale is drawn by the same way as in Figure 1, but its tone changes from the darkest region where $\Delta(B - I)_{\text{DIA}} \leq -0.05$ mag to the brightest region where $\Delta(B - I)_{\text{DIA}} > 0.07$ mag. To better show the fine structures of the map in a region near the caustics, the region enclosed by a box is expanded and presented in the upper panel of Figure 3.

source star, the colour curve of a source transit event is characterized by turns at the moment when the lens enters (or leaves) the source star surface. Measurement of the turning time, $t_\cap$, is important because one can determine the source radius crossing time from the measured value of $t_\cap$ by

$$t_* = \left[\beta^2 + \left(\frac{|t_\cap - t_0|}{t_E} \right)^2 \right]^{1/2} t_E, \quad (13)$$

where the lensing parameters of β , t_0 , and t_E are determined from the overall shape of the light curve. We note that the described patterns of the DIA colour curves are very similar to those of the un-blended event colour curves constructed by using the conventional method (see Han, Park, & Jeong 2000).

The patterns of the colour map for the binary lens system and the resulting colour curves are as follows. First, compared to the single lens map, the binary lens map is much more complex. In the regions around fold caustics, the iso-colour-change contours are parallel with the caustics. However, we note that the contours on the left and right sides of the caustics are not symmetric with respect to the caustic line. In the regions around cusps, on the other hand, the contours form separate peaks of blue colour changes. Second, the colour curves of caustic crossing binary lens events are characterized by the turns during caustic crossings. The colour becomes redder as the source approaches the lens caustics. It becomes most reddish at around the time when the edge of the source star touches the caustics. As the source further approaches to the caustics and thus the inner region of the source star disk lies on the caustics,




Figure 3. The colour map (upper panel) near the caustic region and the colour curves (lower panel) resulting from various source star trajectories (straight lines). The region of the presented map corresponds to the enclosed part of the map in Figure 2. The grey tone of the map, the line types of the source trajectories, and the corresponding colour curves are same as those in Figure 2.

the colour becomes bluer. During each caustic crossing, there are two red peaks which occur when the left and right sides of the source star disk lie on the caustics. By measuring the time separation between the two consecutive red peaks, Δt , one can estimate the approximate value of the source radius crossing time by

$$t_* \sim \frac{\Delta t}{2 \sin \phi}, \quad (14)$$

where ϕ is the angle between the source trajectory and the tangential line of the fold caustic at the position where the source crosses the caustics.[†] The value of ϕ can be determined from the global fit of the binary lens event light curve.

[†] If the fold caustic is a straight line, the relation in equation (14) is exactly valid. However, for an extended source event the

5 UNCERTAINTIES IN DIA COLOUR MEASUREMENTS

In this section, we show by example that although the colour changes induced by microlensing are usually small, they can be measured with uncertainties small enough for one to extract useful information about the lens and source stars. We estimate the uncertainties of measured colours, $\delta[\Delta(m_{\nu_2} - m_{\nu_1})_{\text{DIA}}]$, in the following ways. Since the uncertainty in the determined source star flux in magnitude is related to the signal-to-noise ratio by

$$\delta m_\nu = \frac{(\delta F_{0,\nu}/F_{0,\nu})}{0.4 \ln 10} \sim \frac{1.09}{S/N}, \quad (15)$$

the uncertainty in the measured colour is related to the signal-to-noise ratio by

$$\delta[\Delta(m_{\nu_2} - m_{\nu_1})_{\text{DIA}}] \sim \sqrt{2} \delta m_\nu \sim \frac{1.54}{S/N}. \quad (16)$$

Then, if $S/N \sim 10$, the uncertainty of the colour measurement will be $\delta\Delta(m_{\nu_2} - m_{\nu_1})_{\text{DIA}} \sim 0.15$ mag. The signal measured from the subtracted image is proportional to the source flux variation, i.e. $S \propto (A_\nu - 1)F_{0,\nu}t_{\text{exp}}$, while the noise originates from both the lensed source and blended background stars, i.e. $N \propto [(F_{0,\nu}A_\nu + \langle B \rangle)t_{\text{exp}}]^{1/2}$. Here t_{exp} is the mean exposure time and $\langle B \rangle$ represents the average total flux of faint unresolved stars within the effective seeing disk with a radius θ_{see} . Then the signal-to-noise ratio is computed by

$$S/N = F_{0,\nu}(A_\nu - 1) \left(\frac{t_{\text{exp}}}{F_{0,\nu}A_\nu + \langle B \rangle} \right)^{1/2}. \quad (17)$$

By using equations (16) and (17), we estimate the uncertainties in the measured colours for several example Galactic bulge single lens events with various impact parameters. We assume the events have a common source size of $\theta_* = 0.0756$ and an Einstein time scale of $t_E = 67.5/2$ days by adopting the values of the MACHO Alert 95-30 for which the extended source effect was actually detected (Alcock et al. 1997c). The lensed source is a K-type giant with $I = 14.05$ mag. The observations are assumed to be carried by using a 1 m telescope with a CCD camera that can detect 12 photons/s for $I = 20$ star. Note that the detection rate is determined considering both the efficiency and gain of the CCD camera. Since the major targets for colour measurements are high amplification events with bright source stars, proper exposure time is important to prevent saturation of images. In addition, since the amplification varies rapidly for these events, we leave t_{exp} as a variable so that the measured signal to be $\sim 40,000$ photons, which corresponds to the median flux of the linear sensitivity region for typical modern CCD cameras. We estimate $\langle B \rangle$ by assuming that blended light comes from stars fainter than the crowding limit. Towards the Galactic bulge field, the crowding limit is set when the stellar number density reaches $\sim 10^6$ stars/deg². Based on the model luminosity function constructed by combining those of Holtzman et al. (1998) and Gould, Bahcall, &

relation is an approximation because the fold caustics can no longer be treated as a straight line.




Figure 4. The colour curves of several example microlensing events along with the estimated uncertainties, which are presented by error bars. We assume the events have a common source size of $\theta_* = 0.0756$. For the details of error estimates, see § 5.

Flynn (1997)[‡], this number density corresponds to the de-reddened I -band magnitude of $I_0 \sim 18.1$. The background flux is normalized for stars in the seeing disk with $\theta_{\text{eff}} = 2''.0$.

In Figure 4, we present the colour curves of the events along with the estimated uncertainties, which are presented by error bars. From the figure, one finds that it will be difficult to distinguish colour change curves for different non-source transit events ($\beta > \theta_*$) due to large uncertainties. However, the uncertainties for source transit events ($\beta \leq \theta_*$) are small enough for one to distinguish curves with different values of β and determine the turning times.

Although it is not included in the above uncertainty estimate, another possible source of uncertainty comes from the residual noise due to flat fielding (C. Alard, private communication). Since this noise is constant, it becomes important especially in the wings of the colour change curve at which the signal is very low as $A_\nu - 1 \rightarrow 0$. However, we find that in the region near the peak amplification where most information from the colour curve is obtained, this noise is not important. For example, let us the residual noise δF_ν to be $\sim 5\%$ of the unlensed source star flux, i.e. $\delta F_\nu \sim 0.05 F_{0,\nu}$. In the region close to the source star, e.g. $\beta \lesssim 0.1$ (and thus $A \gtrsim 10$), the uncertainty in the DIA colour measurements estimated by using equations (15) and (16) will be $\lesssim 0.008$ mag for the same source of the example event in the above error estimate. This uncertainty is substantially (nearly an order) smaller than the colour change of $\gtrsim 0.05$ mag for source transit events.

[‡] For the detailed method of the model LF construction, see Han, Jeong, & Kim (1998).

6 SUMMARY

We investigate the chromaticity of single and binary microlensing events when their colour changes are measured by using the recently developed DIA photometric method. The findings from this investigation are summarized as follows:

- (i) The microlensing induced colour change measured by the DIA method differs from that measured by using the conventional photometric method.
- (ii) Despite the difference, from the DIA colour curve one can obtain the same information about the lens and source star as the information obtained from the conventional colour curve. This is because both colour curves depend on the same parameters.
- (iii) However, since the DIA colour curve is not affected by blending, one can determine the lens proper motion and source star surface intensity profile with significantly reduced uncertainties.
- (iv) While the colour of a non-source transit event continues to become redder as the lens approaches the source star, the colour curve of a source transit event is characterized by turns the moments when the lens enters and leaves the source star disk. For the source transit event, one can determine the source radius crossing time by measuring the turning time of the colour curve.
- (v) The colour curves of binary lens events are much less symmetric and vary greatly depending on the lens system geometry and the source star trajectory. For a caustic crossing event, one can measure the source radius crossing time by measuring the time separation between the two consecutive red peaks on the colour curve.

This work was supported by the grant (KRF-99-041-D00442) of the Korea Research Foundation.

REFERENCES

- Alard, C., & Guibert, J. 1997, *A&A*, 326, 1
- Afonso, C., et al. 1998, *A&A*, 337, 17
- Alard, C. 1996, in *Proc. IAU Symp. 173, Astrophysical Applications of Gravitational Lensing*, eds. Kochanek C. S. & Hewitt J. N., (Dordrecht: Kluwer), 215
- Alard, C. 1997, *A&A*, 321, 424
- Alard, C. 1999, *A&A*, 343, 10
- Alard, C., & Guibert, J. 1997, *A&A*, 326, 1
- Alard, C., & Lupton, R. H. 1998, *ApJ*, 503, 325
- Alard, C., Mao, S., & Guibert, J. 1995, *A&A*, 300, L17
- Albrow, M. D., et al. 1999a, *ApJ*, 512, 672
- Albrow, M. D., et al. 1999b, *ApJ*, 522, 1011
- Alcock, C., et al. 1993, *Nature*, 365, 621
- Alcock, C., et al. 1997a, *ApJ*, 479, 119
- Alcock, C., et al. 1997b, *ApJ*, 486, 697
- Alcock, C., et al. 1997c, *ApJ*, 491, 436
- Alcock, C., et al. 1999a, *ApJ*, 518, 44
- Alcock, C., et al. 1999b, *ApJ*, 521, 602
- Alcock, C., et al. 1999c, *ApJS*, 124, 171
- Allen, C. W. 1973, *Astrophysical Quantities*, (London: Athlon Press), 208
- Aubourg, E., et al. 1993, *Nature*, 365, 623
- Buchalter, A., Kamionkowski, M., & Rich, R. M. 1996, *ApJ*, 469, 676
- Di Stefano, R., & Esin, A. A. 1995, *ApJ*, 448, L1
- Gaudi, B. S., & Gould, A. 1999, *ApJ*, 513, 619

- Goldberg, D. M. 1998, *ApJ*, 489, 156
- Goldberg, D. M., & Woźniak, P. R. 1998, *Acta Astron.*, 48, 19
- Gould, A. 1994, *ApJ*, 421, L71
- Gould, A., Bahcall, J., & Flynn, C. 1997, *ApJ*, 482, 913
- Gould, A., & Welch, D. 1996, *ApJ*, 464, 212
- Han, C. 1997, *ApJ*, 490, 51
- Han, C. 1998, *ApJ*, 500, 569
- Han, C. 1999, *MNRAS*, 309, 373
- Han, C., & Kim, T.-W. 1999, *MNRAS*, 305, 795
- Han, C., Jeong, Y., & Kim, H.-I. 1998, *ApJ*, 507, 102
- Han, C., Park, S.-H., & Jeong, J.-H. 2000, *MNRAS*, 316, 97
- Han, C., Park, S.-H., Kim, H.-I., & Chang, K. 2000, *MNRAS*, 316, 572
- Heyrovský, D., & Sasselov, D. 2000, *ApJ*, 529, 69
- Holtzman, J. A., et al. 1998, *AJ*, 115, 1946
- Kamionkowski, M. 1995, *ApJ*, 442, L9
- Loeb, A., & Sasselov, D. 1995, *ApJ*, 449, L33
- Maoz, D., & Gould, A. 1994, *ApJ*, 425, L67
- Nemiroff, R. J., & Wickramasinghe, W. A. D. T. 1994, *ApJ*, 424, L21
- Palanque-Delabrouille, N., et al. 1998, *A&A*, 332, 1
- Peletier, R. F. 1989, *Elliptical Galaxies: Structure and Stellar Content*, PhD Thesis, (Groningen: Kapteyn Institute)
- Sasselov, D. 1997, in *Variable Stars and the Astrophysical Returns of Microlensing Surveys*, eds. R. Ferlet, J.-B. Maillard, & B. Raban (Gif-sur-Yvette: Edition Frontières), 141
- Schmidt-Kaler, Th. 1982, *Landolt Bornstein: Numerical Data and Fundamental relationships in Science and Technology*, eds. Schifers K. & Voigt H. H., (Berlin: Springer)
- Schneider, P., Ehlers, J., & Falco, E. F. 1992, *Gravitational Lenses* (Berlin: Springer), 313
- Tomaney, A. B., & Crotts, A. P. S. 1996, *AJ*, 112, 2872
- Udalski, A., Szymański, M., Kaluźny, J., Kubiak, M., Krzemiński, W., Mateo, M., Preston, G. W., & Paczyński, B. 1993, *Acta Astron.*, 43, 289
- Udalski, A., Kubiak, M., Szymański, M., Pietrzyński, G., Woźniak, P., & Żebruń, K. 1998, *Acta Astron.*, 48, 431
- Valls-Gabaud, D. 1994, in *Large-Scale Structure of the Universe*, eds. J. Mucket et al. (Singapore: World Scientific), 326
- Valls-Gabaud, D. 1998, *MNRAS*, 294, 747
- Van Hamme, W. 1993, *AJ*, 2096
- Witt, H. 1990, *A&A*, 236, 311
- Witt, H., & Mao, S. 1994, *ApJ*, 430, 505
- Woźniak, P., & Paczyński, B. 1997, *ApJ*, 487, 55